

AAP Updates Treatments for Head Lice

4/27/2015

It's probably making you squirm just thinking about it – your child comes home from school scratching his head. He has lice. It's every parent's nightmare, but it is important to remember that head lice is a nuisance, not a serious disease or a sign of poor hygiene.

An updated clinical report by the American Academy of Pediatrics (AAP) "[Head Lice](#)," in the May 2015 issue of *Pediatrics* (published online April 27), provides information to pediatricians and other health practitioners on safe and effective methods for treating head lice, including new products and medications.

Most cases of head lice are acquired outside of school. In the report, the AAP continues to recommend that a healthy child should not be restricted from attending school because of head lice or nits (eggs). Pediatricians are encouraged to educate schools and communities that no-nit policies are unjust and should be abandoned. Children can finish the school day, be treated, and return to school.

Unless resistance has been seen in the community, pediatricians and parents should consider using over-the-counter medications containing 1 percent permethrin or pyrethrins as a first choice of treatment for active lice infestations. The best way to interrupt a chronic lice problem is with regular checks by parents and early treatment with a safe, affordable, over-the-counter pediculicide. After applying the product according to the manufacturer's instructions, parents should follow with nit removal and wet combing. The treatment should be reapplied at day 9, and if needed, at day 18.

In areas with known resistance to an over-the-counter pediculicide, or when parents' efforts on their own do not work, parents should involve their pediatrician for treatment with a prescription medication such as spinosad or topical ivermectin. These are new medications that were introduced since the last time the AAP published recommendations on head lice in 2010.

Once a family member is identified with head lice, all household members should be checked. The AAP does not recommend excessive environmental cleaning, such as home pesticides. However, washing pillow cases and treating natural bristle hair care items that may have been in contact with the hair of anyone found to have head lice are reasonable measures.

While it is unlikely to prevent all cases of head lice, children should be taught not to share personal items such as combs, brushes, and hats. Regular observation by parents can also be an effective way to detect and quickly treat head lice infestations.

###

The American Academy of Pediatrics is an organization of 62,000 primary care pediatricians, pediatric medical subspecialists and pediatric surgical specialists dedicated to the health, safety and well-being of infants, children, adolescents and young adults.

AAP MEDIA CONTACTS

AAP Headquarters

Phone: 847-434-7877

Email: commun@aap.org